CODEN: WJAPAC Impact Factor: 5.48 ISSN: 3049-3013

World Journal of Advance Pharmaceutical Sciences

Volume 3, Issue 11, Page: 19-22 Review Article www.wjaps.com

DRAVYAGUNA VIJNANA AND PHYTOCHEMISTRY: A COMPREHENSIVE REVIEW ON PLANT SECONDARY METABOLITES AND THEIR THERAPEUTIC PHARMACODYNAMICS

Dr. Kishor Kumar Kumatad*[1], Dr. Vidyalaxmi Pujari[2], Dr. Shashidhar Naik[3]

[1]2nd Year PG Scholar, Department of PG Studies in Dravyaguna, BLDE Association's AVS Ayurveda Mahavidyalaya, Hospital & Research Centre, Vijayapura, Karnataka, India.
[2]Associate Professor, Department of PG Studies in Dravyaguna, BLDE Association's AVS Ayurveda Mahavidyalaya, Hospital & Research Centre, Vijayapura, Karnataka, India.
[3]Professor & Head, Department of PG Studies in Dravyaguna, BLDE Association's AVS Ayurveda Mahavidyalaya, Hospital & Research Centre, Vijayapura, Karnataka, India.

How to cite this Article: Dr. Kishor Kumar Kumatad*[1], Dr. Vidyalaxmi Pujari[2], Dr. Shashidhar Naik[3]. (2026). Dravyaguna Vijnana And Phytochemistry: A Comprehensive Review On Plant Secondary Metabolites And Their Therapeutic Pharmacodynamics. World Journal of Advance Pharmaceutical Sciences, 3(11), 19-22.

Copyright © 2026 Dr. Kishor Kumar Kumatad* | World Journal of Advance Pharmaceutical Sciences
This is an open-access article distributed under Creative Commons Attribution-Non Commercial 4.0 International license (CC BY-NC 4.0)

Article Info

Article Received: 12 Sept. 2026,

Article Revised: 02 Oct. 2026,

Article Accepted: 22 Oct. 2026.

***Corresponding author:**

Dr. Kishor Kumar Kumatad 2nd Year PG Scholar, Department of PG Studies in Dravyaguna, BLDE Association's AVS Ayurveda Mahavidyalaya, Hospital & Research Centre, Vijayapura, Karnataka, India.

DOI: https://doi.org/10.5281/zenodo.23850582

ABSTRACT

Plants produce a rich variety of natural compounds known as secondary metabolites. While these chemicals are not required for a plant's basic daily growth, they play vital roles in helping the plant survive, defend itself against pests and diseases, and adapt to harsh environmental conditions. In both modern medicine and traditional Ayurvedic science (Dravyaguna Vijnana), these active plant compounds form the biological foundation for how herbal remedies work and how new medicinal drugs are discovered. This comprehensive review bridges ancient wisdom and modern science by exploring the chemical structures, natural sources, healing mechanisms, and Ayurvedic equivalents of six major groups of plant compounds: Phenolics, Alkaloids, Saponins, Terpenes, Lipids, and Carbohydrates.

KEYWORDS: *Dravyaguna Vijnana, Ayurvedic Pharmacology, Secondary Metabolites, Phytochemistry, Phenolics, Alkaloids, Saponins, Terpenes.*

1. INTRODUCTION

Plant secondary metabolites are organic molecules synthesized by plant cells through secondary metabolic pathways branching off primary metabolic routes (such as the shikimic acid, mevalonic acid, and methylerythritol phosphate pathways). Unlike primary metabolites such as nucleic acids, essential amino acids, and structural polysaccharides secondary metabolites are not directly involved in basic cellular maintenance, growth, or reproduction. However, they are fundamental to plant survival, playing critical roles in defense against herbivory, microbial pathogens (serving as phytoalexins, antibiotics, antifungals, and antivirals), and abiotic stressors such as ultraviolet (UV) radiation through light-absorbing chromophores.

In the domain of Ayurveda and traditional phytomedicine, Dravyaguna Vijnana systematically explores the ultimate pharmacological properties (Rasa, Guna, Veerya, Vipaka, and Prabhava) of Dravya. Modern pharmacological science establishes that these systemic attributes are intrinsically mediated by the chemical architecture and concentration of secondary metabolites contained within plant matrices.

These molecules serve as pivotal lead structures for single-compound drug isolation and standardized phytotherapeutic formulations.

2. MAJOR GROUPS OF SECONDARY METABOLITES

Secondary metabolites possess vast chemical diversity and are systematically grouped into principal structural classes based on their biosynthetic origin, functional groups, and carbon skeletal frameworks:

- **Phenolics:** Compounds characterized by at least one aromatic ring bearing one or more hydroxyl groups.
- **Alkaloids:** Nitrogenous heterocyclic basic natural products with pronounced systemic physiological activity.
- **Saponins:** Amphiphilic glycosides comprising steroidal or triterpenoid aglycones coupled with sugar moieties.
- **Terpenes:** Isoprene-derived (C5H8) volatile and non-volatile polymers exhibiting extensive structural variation.
- **Lipids & Fixed Oils:** Hydrophobic or amphipathic molecules including fatty acid esters, waxes, and essential oils.
- **Therapeutic Carbohydrates:** Complex plant polysaccharides, mucilages, and gums involved in demulcent and bulk action.

3. CHEMICAL ARCHITECTURE & THERAPEUTIC DYNAMICS

3.1 Phenolic Compounds

Phenolics constitute a highly diverse class ranging from simple single-ring structures to complex polymeric assemblies. They originate primarily via the shikimic acid pathway and contribute heavily to floral pigmentation, astringency, and organoleptic properties.

Structural Subclasses of Phenolics

- **Simple Phenolics:** Single aromatic rings substituted with hydroxyl (-OH), carboxyl (-COOH), or aldehyde groups. Examples include Gallic acid (anti-inflammatory, antimicrobial, bronchodilatory), Salicylic acid (analgesic, anti-inflammatory), Capsaicinoids from Capsicum species (circulatory stimulant, rubefacient), and Arbutin from Arctostaphylos uva-ursi (urinary antiseptic). Essential Ayurvedic botanicals rich in simple phenolics include Haridra (Curcuma longa), Tulasi (Ocimum sanctum), and Amlaki (Phyllanthus emblica).
- **Tannins:** Polyphenolic compounds possessing high molecular weight capable of precipitating proteins. Divided into: (a) Hydrolyzable Tannins (gallic or ellagic acid esters with central glucose cores) acting as astringents, antioxidants, and urinary antiseptics; and (b) Condensed Tannins / Proanthocyanidins (flavonoid polymers). Highly rich in Haritaki (Terminalia chebula), Vibhitaki (Terminalia bellirica), Arjuna (Terminalia arjuna), Ashoka (Saraca asoca), and Kutaja (Holarrhena antidysenterica), exerting strong antidiarrheal and hemostatic activities.
- **Coumarins:** Benzo-α-pyrone derivatives (lactones of o-hydroxycinnamic acid) including aesculetin, umbelliferone, and scopoletin. Distributed across Solanaceae, Thymelaeaceae, and Apiaceae. Exhibiting potent anti-inflammatory, anticoagulant, antineoplastic, and neuroprotective properties. Key sources include Ashwagandha (Withania somnifera), Shatavari (Asparagus racemosus), Bhringaraja (Eclipta alba), and Bakuchi (Psoralea corylifolia).
- **Flavonoids:** C6-C3-C6 hydroxylated phenolic structures categorized into flavones, flavonols, and anthocyanins. Celebrated for potent antioxidant, antiallergic, vasoprotective, gastric mucosal protective, and anti-inflammatory effects. Present in Glycyrrhiza glabra (Yastimadhu), Amlaki, and Arjuna.

3.2 Alkaloids

Alkaloids are basic nitrogenous heterocyclic compounds predominantly derived from amino acids (e.g., tryptophan, tyrosine, lysine, ornithine). They are prevalent in specific Angiosperm orders such as Ranunculales, Papaverales, Magnoliales, and Rosales.

Alkaloid basic skeletal types include Acridones, Indoles, Isoquinolines, Piperidines, Pyridines, Pyrrolidines, Quinolines, and Tropanes. Due to their potent target affinity, alkaloids exhibit significant physiological actions: analgesia, local anesthesia, autonomic nerve modulation, cardiac stimulation, vasodilation/constriction, and cytotoxicity against tumor cells.

Prototypical Bioactive & Ayurvedic Alkaloids

- **Nicotine (Nicotiana tabacum):** A potent ganglionic nicotinic acetylcholine receptor agonist exhibiting complex central nervous system and autonomic cardiovascular stimulation.
- **Caffeine (Coffea spp., Camellia sinensis):** A central nervous system xanthine alkaloid stimulant, adenosine receptor antagonist, and renal diuretic.
- **Ayurvedic Alkaloidal Drugs:** Sarpagandha (Rauwolfia serpentina - reserpine), Brahmi (Bacopa monnieri), Mandukaparni (Centella asiatica), Vatsanabha (Aconitum ferox - aconitine), and Bhanga (Cannabis sativa).

3.3 Saponins

Saponins are high-molecular-weight amphiphilic glycosides comprising a lipophilic aglycone (sapogenin) moiety—either a steroid (C27) or a triterpenoid (C30) skeleton—linked to hydrophilic mono- or oligosaccharide chains (pentoses, hexoses, or uronic acids). Their physical property of lowering surface tension produces persistent foam in aqueous solutions and erythrocyte membrane lysis in vitro.

Therapeutically, saponins exhibit notable expectorant, antitussive, anti-inflammatory, adaptogenic, and immunomodulatory actions. For example, Glycyrrhizin from Glycyrrhiza glabra (Yastimadhu) provides potent anti-inflammatory and hepatoprotective effects in chronic hepatitis. Prominent Ayurvedic botanicals containing saponins include Shatavari (Asparagus racemosus), Gokshura (Tribulus terrestris), Bala (Sida cordifolia), and Ashwagandha (Withania somnifera).

3.4 Terpenes and Isoprenoids

Terpenes constitute the largest class of botanical secondary metabolites, synthesized from repeating five-carbon isoprene units (C5H8) via the mevalonate (MVA) or methylerythritol phosphate (MEP) pathways. They are categorized based on unit multiplication:

- **Hemiterpenes (C5 - 1 unit):** Isoamylene, Angelic acid; volatile sensory blocks.
- **Monoterpenes (C10 - 2 units):** Menthol, Camphor, Limonene, Linalool; utilized as counterirritants, topical analgesics, and anthelmintics.
- **Sesquiterpenes (C15 - 3 units):** Farnesol, Caryophyllene, Bisabolol; antimicrobial, anti-inflammatory, and cardiotonic agents.
- **Diterpenes (C20 - 4 units):** Gibberellins, Taxol, Phytol (Vit A/K precursors); antineoplastic, antirheumatic, and diaphoretic dynamics.
- **Sesterterpenes (C25 - 5 units):** Geranylfarnesol derivatives; demonstrating selective cytotoxicity.
- **Triterpenes (C30 - 6 units):** Squalene, Boswellic acids (from Boswellia serrata); profound anti-inflammatory, anti-arthritic, and tissue-remodeling properties.

3.5 Hydrophobic Lipids & Essential Oils

Plant lipids encompass hydrophobic non-polar compounds. While primary lipids serve structural cell membrane roles and energy reserves, specialized lipophilic fractions exhibit powerful therapeutic attributes:

- **Fixed Oils:** Glycerol esters of long-chain unsaturated fatty acids (e.g., Linseed oil, Castor oil, Olive oil). Possess anti-inflammatory PUFA profiles, antioxidant activities, and purgative effects (e.g., Eranda Taila). Rich in Tila (Sesamum indicum) and Mahanimba.
- **Waxes:** Esters of long-chain monohydric alcohols and fatty acids (e.g., Jojoba wax from Simmondsia chinensis), offering cutaneous barrier restoration, wound healing, and anti-inflammatory protection.
- **Essential Oils:** Volatile lipophilic mixtures of monoterpenes, sesquiterpenes, and phenylpropanoids (e.g., Peppermint, Eucalyptus, Clove, Cinnamon oils). Function as powerful antimicrobials, antispasmodics, analgesics, and carminatives.

3.6 Therapeutic Carbohydrates and Polysaccharides

Although fundamental primary metabolic products (carbon and energy storage), complex carbohydrates, plant gums, and mucilages act as key therapeutic agents. Mucilages (found in Flaxseed, Isabgol) absorb water to form protective viscous gels, exerting demulcent, mucosal soothing, and anti-ulcer actions. Plant gums (Acacia, Tragacanth) serve as bulk laxatives and pharmaceutical excipients.

4. PHYTOCHEMICAL SYNTHESIS & AYURVEDIC MAPPING

The integrated comparative matrix below maps secondary metabolite classes to their chemical characteristics, representative bioactive examples, primary modern therapeutic activities, and classic Ayurvedic phytomedicines rich in these metabolites.

Class	Chemical Characteristics	Representative Examples	Therapeutic Actions	Ayurvedic Drugs (Dravya)
Phenolics	Aromatic ring with one or more -OH groups; range from simple monomers to polymers.	Gallic acid, Salicylic acid, Quercetin, Tannins, Coumarins	Anti-inflammatory, Antioxidant, Antimicrobial, Vasoprotective, Astringent	Haridra, Tulasi, Amlaki, Arjuna, Ashoka, Haritaki
Alkaloids	Nitrogenous heterocyclic basic compounds; derived from amino acids.	Nicotine, Caffeine, Reserpine, Morphine, Aconitine	Analgesic, CNS stimulation/depression, Antihypertensive, Antineoplastic	Sarpagandha, Brahmi, Mandukaparni, Vatsanabha, Bhanga
Saponins	Amphiphilic glycosides; steroidal or triterpenoid aglycone + sugar chain.	Glycyrrhizin, Diosgenin, Ginsenosides	Expectorant, Adaptogenic, Immunomodulatory, Hepatoprotective, Hemolytic	Shatavari, Gokshura, Yastimadhu, Ashwagandha, Bala
Terpenes	Isoprene units (C5H8)n; volatile and non-volatile structures.	Menthol, Camphor, Caryophyllene, Boswellic acids	Analgesic, Anti-inflammatory, Anti-arthritic, Anthelmintic, Antimicrobial	Haridra, Tulasi, Bhringaraja, Gokshura, Shallaki (Boswellia)
Lipids & Oils	Hydrophobic esters, triglycerides, fatty acids, and	Linseed oil, Castor oil, Cineole, Eugenol,	Anti-inflammatory, Laxative/Purgative,	Tila, Eranda, Musta, Mahanimba, Devadaru

	volatile terpene mixtures.	Jojoba wax	Cutaneous healing, Antimicrobial	
Carbohydrates	Polysaccharides, mucilages, and gums (C, H, O polymers).	Mucilage, Acacia gum, Tragacanth, Cellulose	Demulcent, Anti-ulcer, Gastroprotective, Bulk laxative	Isabgol, Flaxseed (Atasi), Marshmallow, Shalmali
Coumarins & Flavonoids	Benzo-α-pyrones & C6-C3-C6 polyphenolic structures.	Umbelliferone, Scopoletin, Rutin, Kaempferol	Anticoagulant, Vasoprotective, Anti-Alzheimer's, Gastric mucosa protection	Bakuchi, Bhringaraja, Shatavari, Neem, Arjuna

5. DISCUSSION & AYURVEDIC INTEGRATION

The detailed understanding of secondary metabolites provides a robust biological validation for the empirical concepts described in Dravyaguna Vijnana. In Ayurvedic pharmacology, the action of a crude drug is attributed to its multi-constituent harmony (Synergy or Polypharmacology), preventing single-target toxicity while providing broad therapeutic efficacy. For instance, the presence of tannins and flavonoids in Terminalia arjuna explains its dual astringent (Kashaya Rasa) and cardioprotective (Hridya) action. Similarly, saponins in Asparagus racemosus validate its Rasayana (rejuvenative) and mucosal-soothing properties.

Contemporary pharmaceutical research increasingly focuses on standardization, bio-guided fractionation, and the identification of lead compounds from medicinal herbs. Understanding secondary metabolites enables Ayurvedic scholars to standardize traditional formulations (e.g., Kwatha, Churna, Taila, Asava-Arishta), ensure batch-to-batch consistency, and establish safety profiles in line with global pharmacopoeial standards.

6. CONCLUSION

Plant secondary metabolites represent an indispensable reservoir of bioactive chemicals that bridge ancient Ayurvedic therapeutics with modern molecular pharmacology. A scientific comprehension of Phenolics, Alkaloids, Saponins, Terpenes, Lipids, and Carbohydrates enriches Dravyaguna Vijnana, facilitating drug discovery, quality control, and evidence-based global acceptance of Ayurvedic phytomedicines.

7. REFERENCES

1. Evans WC. Trease and Evans' Pharmacognosy. 16th ed. Saunders Elsevier, 2009.
2. Harborne JB. Phytochemical Methods: A Guide to Modern Techniques of Plant Analysis. 3rd ed. Chapman & Hall, 1998.
3. Bruneton J. Pharmacognosy, Phytochemistry, Medicinal Plants. 2nd ed. Intercept Ltd, 1999.
4. Sharma PV. Dravyaguna-Vijnana. Vol. I & II. Chaukhambha Bharati Academy, 2006.
5. Wink M. Annual Plant Reviews, Volume 39: Functions and Biotechnology of Plant Secondary Metabolites. 2nd ed. Wiley-Blackwell, 2010.
6. World Health Organization. Quality Control Methods for Medicinal Plant Materials. World Health Organization, 2011.